
From: jeffrey E. <jeevacation@gmail.com>
Sent: Wednesday, July 20, 2016 10:55 AM
To: Eva Dubin

Not recommended
for treating
malaria in children
who weigh less
than 11
kgs.

For children who weigh less than
11 kgs talk to your doctor. There

.

If you forgot to
take
a dose,
don't worry. Just
take
your
next dose as
soon as
you remember. Then continue
your
treatment as before.

Don't take a double dose to make up for
a missed dose

.

Don't stop Malarone without advice

Keep
taking

Malarone for
7 days after you return to a malaria

.

For
area

.

Take
the full course of
Malarone
for
maximum protection. Stopping
early puts
you
at risk of getting malaria, as it
takes 7 days to ensure that any parasites that may be in your blood following a bite from an infected mosquito are killed.

Do not

tak

e Malarone for

longer than 37 days. If you are expecting to stay in an area with malaria for more than 28 days check with your doctor before travelling. Malarone may not be suitable for you.

5

)

Possib

I

e side effects

Like all medicines, Malarone can cause side effects, although not everybody gets them.

Look out for the following severe

reactions. They have occurred in a small number of people, but their exact frequency is unknown.

Severe allergic reactions

-

signs include:

.

rash and itching

.

sudden wheezing, tightness of the

chest or throat, or difficulty breathing = swollen eyelids, face, lips, tongue or other part of the body.

Contact a doctor immediately

if you get any of these

s

ymptoms.

Stop

taking Malarone

.

Severe skin reactions

.

skin rash which may blister and looks like small targets (central dark spots, surrounded by paler area with a dark ring around the edge) (erythema multiforme

)

.

severe widespread rash with blisters

and peeling skin, particularly occurring around the mouth, nose, eyes and genitals (Stevens Johnson syndrome

).

If you notice any of these symptoms

contact a doctor urgently.

Most of the other side effects reported

have been mild and have not lasted

very

long.

Very common side effects

These may affect

more than 1 in

10 people:

.

headache

•
feeling sick and being sick (

nausea

and

vomiting

=

•
stomach pa

in

•
diarrhoea.

Common side effects

These may affect

up to 1 in 10 people:

•
(neutropenia

) which may make you
more likely to catch infectio=s

•
low levels of sod=um in the blood

(hyponatraem

=a)

•</=iv>an increase in liver enzy

--

=A0 please note

The information contained in this commu=ication is confidential, may be attorney-client privileged, may cons=titute inside information, and is intended only for the use of the addre=see. It is the property of JEE Unauthorized use, disclosure or copyi=g of this communication or any part thereof is strictly prohibited a=d may be unlawful. If you have received this communication in error, pl=ase notify us immediately by return e-mail or by e-mail to jeevacation@gmail.com, a=d destroy this communication and all copies thereof, including all a=tachments. copyright -all rights reserved