
From: [REDACTED] >
Sent: Thursday, November 14, 2013 1:54 AM
To: Jeffrey Epstein
Subject: Isoprostane in urine could be assay

In absence of peripheral type 2 dm there should be no excuse for elevated le=els excreted in urine. Only other place is brain. The oxidative stress stor= is stupid and what is likely happening is metabolic strain. Can't absorb (=nsulin resistance neurons) rather than trying hard to do something. It woul= also be quantitative.

You'd see it early

Not sure anything w acetylcholine would be useful unless in CSF and that's t=oo invasive.

Blood assays not useful, since you'd have to draw from the internal Jugular t= be sure it even came from brain metabolism.=?xml version=.0" encoding=TF-8"?> <!DOCTYPE plist PUBLIC "-//Apple//DTD PLIST 1.0//EN"
"http://www.apple.com/DTDs/PropertyList-1.0.dtd">
<plist version=.0">
<dict>
 <key>conversation-id</key>
 <integer>273002</integer>
 <key>date-last-viewed</key>
 <integer>0</integer>
 <key>date-received</key>
 <integer>1384394071</integer>
 <key>flags</key>
 <integer>8590195713</integer>
 <key>gmail-label-ids</key>
 <array>
 <integer>6</integer>
 <integer>2</integer>
 </array>
 <key>remote-id</key>
 <string>361808</string>
</dict>
</plist>